

## (別紙1) 裸眼視力1.0未満の幼児・児童・生徒の割合

(単位：%)

	幼稚園		小学校		中学校		高等学校	
	佐賀	全国	佐賀	全国	佐賀	全国	佐賀	全国
H1	10.4	25.8	20.6	20.6	37.5	40.9	53.3	55.8
H2	4.7	18.6	20.5	21.2	41.2	41.6	55.1	56.4
H3	11.5	21.6	21.3	21.7	39.9	43.6	56.2	57.5
H4	11.6	20.2	22.0	22.5	42.9	45.6	58.0	59.2
H5	10.9	20.6	23.8	23.8	45.1	47.3	60.4	61.9
H6	10.4	23.8	24.9	24.7	48.9	48.8	62.1	62.3
H7	16.1	26.8	26.0	25.4	48.6	49.1	62.0	61.8
H8	25.8	21.4	26.6	25.8	50.4	49.8	62.7	62.7
H9	20.4	23.1	27.1	26.3	51.2	49.7	63.5	63.2
H10	5.8	25.8	27.2	26.3	51.9	50.3	64.6	62.5
H11	11.7	24.0	25.8	25.8	52.3	49.7	66.8	63.3
H12	2.4	28.7	26.4	25.3	51.3	50.0	64.7	62.5
H13	31.5	27.2	26.3	25.4	51.1	48.2	65.7	60.3
H14	13.3	26.5	25.6	25.7	49.7	49.0	67.3	63.8
H15	35.3	25.3	26.2	25.6	49.6	47.8	64.0	60.0
H16	10.4	20.8	27.7	25.6	50.5	47.7	61.6	59.3
H17	8.3	20.4	27.1	26.5	49.2	47.8	63.5	58.4
H18	12.0	24.1	29.2	28.4	49.3	50.1	62.7	58.7
H19	12.0	26.2	28.1	28.1	53.8	51.2	58.6	55.4
H20	X	<u>28.9</u>	30.8	29.9	53.1	52.6	68.7	58.0
H21	X	24.9	30.9	29.7	52.6	52.5	60.7	59.4
H22	X	26.4	31.9	29.9	55.9	52.7	64.8	55.6
H23	X	25.5	32.0	29.9	49.6	51.6	53.5	60.9
H24	X	27.5	32.7	30.7	56.3	54.4	64.1	64.5
H25	X	24.5	30.9	30.5	48.8	52.8	X	65.8
H26	X	26.5	31.9	30.2	51.1	53.0	53.4	62.9
H27	X	26.8	33.4	31.0	52.0	54.1	X	63.8
H28	X	27.9	33.6	31.5	55.2	54.6	X	66.0
H29	X	24.5	34.1	32.5	53.6	56.3	X	62.3
H30	X	26.7	38.1	34.1	52.3	56.0	63.0	67.2
R1	X	26.1	37.8	34.6	57.6	57.5	70.6	67.6
R2	X	27.9	38.7	37.5	54.9	58.3	X	63.2
R3	X	24.8	39.0	36.9	66.3	60.7	X	70.8
R4	X	25.0	<u>40.5</u>	<u>37.9</u>	X	<u>61.2</u>	68.7	<u>71.6</u>
R5	X	22.9	39.4	37.8	62.0	60.9	X	67.8

注1) 「X」は、疾病・異常被患率等の標準誤差が5以上、受検者数が100人(5歳は50人)未満、  
回答校が1校以下又は疾病・異常被患率が100%のため統計数値を公表しない。

注2) 下線の部分は、平成元年度以降最高を示す。

注3) 令和2年度から5年度は新型コロナウイルス感染症の影響で、他の年度と測定時期が異なっているため、  
過去の数値と単純比較することはできない。

注4) 昨年度示したもので過去の数値に一部誤りがあり訂正している。

(H18の小学校・中学校・高等学校(佐賀)、H30の高等学校(全国))

## (別紙2) 耳疾患の被患率

(単位：%)

	幼稚園		小学校		中学校		高等学校	
	佐賀	全国	佐賀	全国	佐賀	全国	佐賀	全国
H7	4.1	1.9	5.3	3.4	3.6	1.7	2.5	0.9
H8	2.2	1.7	5.9	3.6	4.6	1.9	2.2	0.9
H9	3.5	1.9	6.1	3.8	3.7	2.0	2.3	1.0
H10	3.3	1.5	5.8	3.7	3.2	2.1	1.3	0.8
H11	2.6	1.8	6.0	4.2	3.1	2.3	2.1	1.1
H12	2.1	1.9	4.0	4.1	3.0	2.5	1.9	1.2
H13	4.3	2.1	5.7	4.2	3.2	2.4	2.1	1.3
H14	<u>8.5</u>	2.0	5.3	4.3	2.8	2.6	1.4	1.7
H15	0.4	2.2	5.6	4.5	2.8	2.7	1.7	1.2
H16	3.2	1.8	4.6	4.3	2.5	2.7	1.9	1.3
H17	3.3	2.1	4.6	4.5	3.1	2.8	2.0	1.3
H18	6.3	2.9	5.4	4.9	3.6	3.1	<u>4.3</u>	1.7
H19	3.7	2.6	5.4	5.1	4.6	3.3	2.1	1.7
H20	1.7	2.8	5.6	5.2	<u>5.5</u>	3.6	2.6	2.0
H21	5.4	2.9	6.3	5.5	4.0	3.4	1.8	2.0
H22	5.4	<u>3.3</u>	6.0	5.4	4.9	3.6	1.8	1.6
H23	4.9	2.5	5.8	5.5	4.0	3.3	2.3	1.6
H24	3.4	2.6	6.0	5.4	3.4	3.6	1.9	1.9
H25	5.5	2.6	7.1	5.4	4.3	3.9	2.4	2.2
H26	2.1	2.3	6.2	5.7	4.5	4.0	2.2	2.1
H27	1.8	2.2	6.4	5.5	4.8	3.6	1.8	2.0
H28	1.9	2.8	6.3	6.1	3.6	4.5	2.3	2.3
H29	1.5	2.3	6.5	6.2	3.5	4.5	2.5	2.6
H30	3.8	2.3	6.0	6.5	4.9	4.7	2.3	2.5
R1	2.3	2.6	7.0	6.3	4.9	4.7	1.9	<u>2.9</u>
R2	2.0	2.0	<u>8.2</u>	6.1	4.7	<u>5.0</u>	2.2	2.5
R3	1.1	2.0	6.6	<u>6.8</u>	<u>5.5</u>	4.9	1.9	2.5
R4	2.1	2.4	5.9	6.6	5.3	4.8	2.5	2.3
R5	1.7	2.2	6.2	6.3	4.8	4.9	2.3	2.6

注1) 下線の部分は、調査実施以来最高を示す。

注2) 令和2年度から5年度は新型コロナウイルス感染症の影響で、他の年度と測定時期が異なっているため、過去の数値と単純比較することはできない。

注3) 昨年度示したもので過去の数値に一部誤りがあり訂正している。

(H18の小学校・中学校・高等学校(佐賀))

## (別紙3) 鼻・副鼻腔疾患の被患率

(単位：%)

	幼稚園		小学校		中学校		高等学校	
	佐賀	全国	佐賀	全国	佐賀	全国	佐賀	全国
H7	4.6	3.3	13.9	<u>9.9</u>	13.9	8.1	10.8	<u>5.8</u>
H8	3.6	2.7	14.0	10.0	14.5	<u>7.8</u>	11.8	5.9
H9	6.8	2.7	15.9	10.0	13.0	8.9	11.3	6.5
H10	4.7	2.9	15.3	10.2	13.1	8.4	12.0	<u>5.8</u>
H11	5.8	2.9	16.4	10.4	12.4	9.0	9.3	6.8
H12	3.9	3.3	13.8	11.3	11.0	9.4	9.9	7.1
H13	6.5	3.2	15.7	10.7	10.4	9.0	10.6	7.4
H14	6.5	3.5	13.7	11.2	11.3	9.4	10.5	7.7
H15	0.6	3.0	13.4	11.1	10.8	10.1	10.1	7.4
H16	4.2	2.7	11.7	10.4	9.7	9.3	8.6	6.8
H17	3.8	3.2	11.2	11.2	11.4	10.6	15.2	8.1
H18	6.9	3.4	14.1	11.9	12.7	10.7	15.8	8.2
H19	2.7	3.7	14.1	12.0	13.6	11.1	11.3	8.4
H20	2.1	3.8	11.5	11.9	12.0	10.8	11.8	8.8
H21	X	4.0	12.8	12.6	12.4	10.8	18.2	9.6
H22	2.0	3.4	14.4	11.7	12.9	10.7	12.7	8.5
H23	10.3	4.4	12.1	12.5	11.9	11.8	14.2	8.8
H24	X	3.5	12.5	12.2	13.8	11.4	14.3	8.6
H25	2.7	3.4	12.6	12.1	13.2	11.1	13.3	8.7
H26	2.4	3.1	13.7	12.3	13.3	11.2	13.8	8.7
H27	1.8	3.6	11.9	11.9	12.3	10.6	8.0	7.3
H28	8.9	3.6	10.9	12.9	10.8	11.5	11.1	9.4
H29	2.0	2.9	12.7	12.8	10.6	11.3	13.0	8.6
H30	3.8	2.9	13.7	13.0	13.1	11.0	9.7	9.9
R1	4.2	3.2	12.6	11.8	12.1	12.1	10.2	9.9
R2	4.8	<u>2.4</u>	11.0	11.0	9.1	10.2	7.9	6.9
R3	2.9	3.0	9.7	11.9	9.3	10.1	8.5	8.8
R4	2.0	3.0	<u>8.3</u>	11.4	<u>8.3</u>	10.7	6.4	8.5
R5	2.6	3.0	10.2	12.4	8.6	10.5	<u>6.1</u>	7.6

注1) 「X」は、疾病・異常被患率等の標準誤差が5以上、受検者数が100人（5歳は50人）未満、回答校が1校以下又は疾病・異常被患率が100%のため統計数値を公表しない。

注3) 昨年度示したもので過去の数値に一部誤りがあり訂正している。

注3) 令和2年度から5年度は新型コロナウイルス感染症の影響で、他の年度と測定時期が異なっているため、過去の数値と単純比較することはできない。

注4) 昨年度示したもので過去の数値に一部誤りがあり訂正している。

(H18の小学校・中学校・高等学校(佐賀))

## (別紙4) むし歯(う歯)の被患率

(単位：%)

	幼稚園		小学校		中学校		高等学校	
	佐賀	全国	佐賀	全国	佐賀	全国	佐賀	全国
H1	80.1	80.9	94.0	90.3	94.9	90.4	96.8	94.2
H2	87.0	80.4	95.3	89.5	95.0	90.0	98.1	93.7
H3	86.3	80.8	94.4	89.3	93.3	89.6	96.4	93.0
H4	87.7	78.7	94.5	89.1	92.9	88.9	96.8	92.6
H5	82.8	75.7	94.3	88.4	90.7	87.8	96.5	91.3
H6	82.0	77.0	93.6	88.0	92.3	87.7	96.8	92.0
H7	80.4	74.7	94.1	87.3	90.6	86.6	95.0	90.6
H8	77.5	73.7	92.5	85.7	87.7	84.8	94.4	90.1
H9	82.0	71.2	91.0	84.7	87.7	83.7	95.9	89.4
H10	75.8	67.7	88.4	82.1	87.3	81.9	94.2	88.2
H11	76.6	67.0	88.9	80.8	85.6	80.1	93.9	86.5
H12	71.7	64.4	85.8	77.9	82.5	76.9	90.7	85.0
H13	74.4	61.5	84.4	75.6	80.3	73.8	91.5	83.7
H14	72.7	61.5	81.7	73.9	76.4	71.2	90.0	82.3
H15	76.7	58.8	79.8	71.3	74.1	67.7	88.7	77.9
H16	66.0	56.9	79.4	70.4	72.4	64.6	85.0	76.0
H17	71.6	54.4	73.2	68.2	66.7	62.7	81.7	72.8
H18	61.5	55.2	74.0	67.8	67.2	59.7	80.8	70.1
H19	62.6	53.7	71.2	65.5	62.7	58.1	75.3	68.5
H20	59.9	50.3	70.8	63.8	56.5	56.0	66.9	65.5
H21	59.4	46.5	68.6	61.8	50.7	52.9	72.9	62.2
H22	59.3	46.1	64.5	59.6	48.1	50.6	69.5	60.0
H23	54.3	43.0	62.9	57.2	46.6	48.3	64.5	58.5
H24	57.3	42.9	59.9	55.8	44.0	45.7	64.1	57.6
H25	52.4	39.5	60.5	54.1	39.8	44.6	62.7	55.1
H26	45.9	38.5	59.8	52.5	35.1	42.4	55.3	53.1
H27	46.5	36.2	58.3	50.8	36.0	40.5	55.0	52.5
H28	47.7	35.6	56.2	48.9	34.9	37.5	53.7	49.2
H29	48.4	35.5	54.5	47.1	35.1	37.3	50.0	47.3
H30	44.3	35.1	53.1	45.3	37.3	35.4	48.3	45.4
R1	X	31.2	51.8	44.8	30.9	34.0	46.7	43.7
R2	38.9	30.3	47.8	40.2	27.4	32.2	44.5	41.7
R3	X	26.5	49.9	39.0	28.0	30.4	43.1	39.8
R4	36.0	24.9	46.2	37.0	28.8	28.2	<u>41.6</u>	38.3
R5	28.2	<u>22.6</u>	<u>42.9</u>	<u>34.8</u>	<u>26.7</u>	<u>28.0</u>	42.2	<u>36.4</u>

注1) 「X」は、疾病・異常被患率等の標準誤差が5以上、受検者数が100人(5歳は50人)未満、  
回答校が1校以下又は疾病・異常被患率が100%のため統計数値を公表しない。

注2) 二重下線の部分は、調査実施以来最低を示す。

注3) 令和2年度から5年度は新型コロナウイルス感染症の影響で、他の年度と測定時期が異なっているため、  
過去の数値と単純比較することはできない。

注4) 昨年度示したもので過去の数値に一部誤りがあり訂正している。

(H18の小学校・中学校・高等学校(佐賀))

## (別紙5) 年齢別身長平均値の推移 (佐賀県)

男子

(単位 : cm)

区分	5歳	6	7	8	9	10	11	12	13	14	15	16	17
S 23	105.0	107.7	112.3	116.7	121.1	125.6	129.6	134.3	136.5	143.3	153.0	156.0	159.6
43	107.9	113.4	118.9	124.1	128.9	133.3	138.4	144.5	151.7	158.1	163.1	165.7	166.5
63	110.5	116.7	121.7	127.5	132.4	137.6	142.8	150.0	157.4	163.4	167.2	168.7	170.0
H 1	110.6	116.0	121.8	127.3	132.9	138.4	143.8	150.4	157.5	163.9	167.3	169.0	170.6
2	110.7	116.4	122.4	127.4	132.7	137.8	143.4	149.9	157.8	163.7	167.8	169.0	170.1
3	110.6	117.0	122.2	127.4	132.3	137.8	144.0	150.9	157.7	164.1	167.8	169.0	169.9
4	110.8	116.6	122.4	127.6	132.5	138.0	143.8	150.8	158.6	164.2	167.7	169.5	170.2
5	110.9	116.5	121.9	127.8	132.9	138.7	144.3	150.6	158.2	164.1	167.4	169.3	170.2
6	110.6	116.4	122.6	127.6	133.2	138.9	144.5	151.0	158.6	163.9	166.9	169.9	169.8
7	110.6	116.6	121.7	127.5	133.0	138.0	144.3	151.0	158.7	164.5	168.0	169.3	170.1
8	110.5	116.8	122.5	127.6	133.1	137.7	144.6	151.4	158.0	164.4	167.8	169.3	169.7
9	110.1	116.1	122.2	127.5	133.2	138.8	144.3	151.8	159.0	164.7	168.3	169.3	170.5
10	110.2	116.6	122.3	127.9	132.8	138.4	144.9	151.9	159.2	164.5	168.0	169.8	170.4
11	110.5	116.4	122.2	128.2	133.1	138.4	144.6	152.5	159.2	165.0	167.4	169.4	170.0
12	110.3	116.3	122.3	127.8	132.8	138.7	144.8	151.8	159.2	164.7	168.4	169.9	170.4
13	110.5	116.7	122.2	127.9	133.2	138.6	144.3	152.5	159.4	164.9	167.7	170.0	170.5
14	110.4	116.7	122.6	128.1	132.6	138.7	145.0	151.5	159.5	164.8	167.7	169.7	170.9
15	110.8	116.4	122.1	127.8	133.3	138.5	144.5	151.8	159.2	165.1	168.0	170.0	170.5
16	110.5	116.7	122.4	128.0	133.4	138.6	144.9	151.9	159.4	164.8	168.5	169.6	170.1
17	110.6	116.5	122.4	127.7	133.1	138.8	144.8	150.8	159.1	164.4	167.7	169.4	170.2
18	110.5	116.4	122.4	128.0	133.5	138.3	144.5	151.9	159.2	164.6	168.1	169.2	170.6
19	110.6	116.5	122.5	128.3	133.6	138.9	144.8	152.0	159.3	164.6	167.8	169.5	171.2
20	110.6	116.8	122.4	127.9	133.4	138.8	145.1	151.8	159.3	165.1	167.6	169.4	170.2
21	110.4	116.1	122.4	127.8	132.9	138.4	144.4	152.0	159.2	164.5	167.7	169.2	169.8
22	109.9	116.3	122.6	128.1	133.2	138.7	145.0	151.7	158.7	164.8	167.6	169.1	170.6
23	110.1	116.2	121.8	128.3	133.2	138.4	144.6	151.8	159.4	164.7	167.8	169.3	170.7
24	110.4	116.3	122.3	128.3	133.8	139.1	144.6	151.5	159.1	164.6	168.3	169.4	170.3
25	110.2	116.7	122.2	128.0	133.4	138.4	144.3	151.7	159.2	164.8	168.2	168.8	170.5
26	110.2	116.5	122.3	127.6	133.6	138.3	144.9	152.3	159.3	164.8	167.7	169.6	170.1
27	110.1	116.5	122.1	127.6	132.7	138.7	144.9	152.5	159.3	165.2	167.9	169.3	171.0
28	109.7	116.3	122.3	127.7	133.3	138.6	144.6	152.7	159.5	165.2	167.7	170.2	170.7
29	110.1	116.2	122.3	128.2	133.2	138.7	145.3	151.7	159.9	164.8	168.0	169.7	171.0
30	110.3	116.4	122.1	127.6	133.5	137.9	144.9	152.1	159.3	164.9	168.1	169.3	170.1
R 1	110.0	116.4	122.3	127.8	133.3	138.6	145.2	152.2	159.2	164.9	167.5	169.6	170.2
2	111.5	116.6	122.7	128.2	133.0	139.1	145.2	151.9	159.7	165.1	167.5	168.3	169.5
3	110.8	116.5	122.2	128.2	133.7	139.0	144.9	152.8	160.0	165.4	167.7	168.9	170.8
4	111.0	116.9	122.2	128.2	133.5	139.5	145.3	153.4	160.8	165.1	168.4	169.2	170.1
5	110.3	116.8	122.9	128.3	133.4	139.3	145.7	153.2	160.7	165.6	168.3	170.1	170.9

女子

(単位 : cm)

区分	5歳	6	7	8	9	10	11	12	13	14	15	16	17
S 23	104.8	106.9	111.5	115.9	120.5	124.8	129.5	135.4	139.7	144.7	150.3	151.7	152.8
43	107.0	112.2	117.9	123.3	128.5	134.5	140.7	146.5	150.5	153.0	154.0	154.2	154.7
63	109.7	115.5	121.2	126.9	132.5	139.4	145.1	150.5	154.3	155.8	156.7	157.5	157.8
H 1	109.6	115.8	121.2	127.2	132.4	139.3	146.1	151.2	154.5	156.2	156.5	157.3	157.6
2	109.6	115.5	121.7	127.2	133.3	139.0	146.2	151.1	154.3	156.1	157.0	157.2	157.7
3	110.0	115.8	121.0	127.1	132.8	139.1	145.7	151.2	154.5	156.2	157.0	158.0	157.2
4	109.9	115.5	121.3	127.3	132.6	139.5	146.4	151.1	154.5	156.3	156.8	157.5	157.6
5	109.5	115.7	121.4	127.2	133.2	140.2	145.9	151.1	154.2	156.3	156.9	157.9	157.9
6	109.7	115.8	121.3	127.3	132.6	140.0	146.6	151.2	154.2	156.3	156.9	157.0	157.3
7	110.0	115.6	121.3	126.9	133.2	139.7	146.3	151.3	154.7	156.4	157.1	157.4	157.6
8	109.8	115.5	120.9	127.1	133.7	139.5	145.8	151.6	154.7	156.4	157.2	157.9	157.9
9	109.3	115.5	121.1	127.2	133.2	140.4	146.8	151.7	154.9	156.1	157.4	158.0	158.1
10	109.7	115.9	121.5	127.2	133.3	140.3	147.1	151.9	154.7	156.0	156.7	157.6	157.9
11	109.8	115.3	121.4	127.2	132.9	139.7	147.3	151.6	154.8	156.0	156.5	157.7	157.3
12	109.5	115.8	121.4	127.3	133.3	140.2	146.6	152.1	154.9	156.4	156.7	157.2	157.7
13	109.8	116.0	121.4	127.4	133.6	139.9	146.2	151.4	155.4	156.4	156.2	157.1	157.7
14	109.9	115.5	121.9	127.3	133.4	140.4	147.1	151.6	155.0	156.2	156.7	157.4	157.7
15	109.9	115.7	121.5	127.4	133.3	140.1	146.8	151.7	154.7	156.4	156.5	156.8	157.3
16	110.0	115.8	121.7	127.6	133.2	140.5	146.4	151.5	154.5	156.4	156.5	157.4	157.3
17	109.6	116.0	121.9	127.2	133.6	140.4	147.6	151.6	154.9	156.3	156.6	157.3	158.1
18	109.6	115.6	121.3	127.3	133.2	139.7	147.2	151.7	154.9	156.6	157.4	157.4	157.4
19	109.3	115.3	121.8	127.1	133.4	140.2	147.6	151.5	155.1	156.1	156.6	157.9	156.8
20	109.9	115.7	120.9	127.3	133.9	140.6	146.8	151.7	155.0	156.3	157.2	157.4	157.7
21	109.9	115.4	121.3	126.9	133.9	140.3	146.8	152.1	154.9	156.2	156.9	157.2	157.3
22	109.1	115.9	121.4	127.0	133.5	140.2	146.8	151.9	154.4	156.4	157.0	157.2	157.2
23	109.6	115.6	121.3	126.7	133.7	140.1	147.3	151.2	155.0	156.2	157.0	157.4	158.0
24	109.7	115.6	121.2	126.5	133.3	140.4	146.7	152.0	154.2	156.4	157.1	157.4	157.6
25	109.5	115.1	121.5	126.8	133.2	139.9	147.0	151.6	154.6	156.2	156.9	157.5	157.1
26	109.6	115.6	121.1	127.4	133.5	140.6	146.5	151.8	154.4	156.1	156.4	157.8	157.5
27	109.2	115.9	121.2	127.0	133.5	140.0	146.9	151.6	154.5	156.2	156.5	157.0	157.4
28	108.8	115.2	121.2	127.1	132.9	140.0	146.8	151.7	154.8	156.4	156.6	157.0	157.0
29	109.5	115.6	120.8	127.3	133.3	139.5	147.0	151.6	154.2	156.2	156.1	157.3	157.5
30	108.8	115.3	120.9	126.8	133.5	139.9	146.4	151.3	154.5	155.9	156.6	157.7	157.5
R 1	109.5	115.9	121.3	127.3	133.1	140.2	146.6	151.5	154.4	155.8	156.4	157.1	157.1
2	110.5	116.1	121.2	127.5	133.5	139.8	147.3	151.7	154.4	155.8	156.3	157.9	156.5
3	109.8	115.5	122.2	127.5	133.7	140.4	147.0	151.5	154.4	155.9	157.3	157.8	157.4
4	109.4	115.8	121.7	127.5	134.0	140.7	147.2	151.7	154.2	156.1	156.6	157.0	157.3
5	109.8	115.6	121.8	127.8	134.7	141.5	147.8	152.0	154.7	155.9	157.4	157.0	157.3

注1) 下線の部分は、調査実施以来最高を示す。

注2) 令和2年度から5年度は新型コロナウイルス感染症の影響で、他の年度と測定時期が異なっているため、過去の数値と単純比較することはできない。

(別紙6) 年齢別体重平均値の推移 (佐賀県)

男子

(単位 : kg)

区分	5歳	6	7	8	9	10	11	12	13	14	15	16	17
S 23	16.4	18.4	20.1	21.7	23.6	25.6	28.2	30.7	31.7	33.0	43.7	47.6	50.3
4 3	17.9	19.4	21.7	23.8	26.3	28.8	31.7	35.9	41.3	46.7	51.6	54.9	56.5
6 3	19.1	21.1	23.2	26.3	29.0	32.4	35.8	41.3	46.8	52.2	57.5	58.8	60.5
H 1	19.1	21.0	23.6	26.1	29.6	33.2	36.9	41.5	46.8	52.4	57.2	58.8	61.5
2	19.3	21.3	23.9	26.5	29.3	32.6	36.5	41.8	47.2	52.7	58.5	60.4	60.9
3	19.3	21.5	23.8	26.8	29.4	32.6	36.8	43.0	48.2	52.9	58.1	60.4	61.6
4	19.3	21.3	24.0	26.5	29.4	33.1	37.1	42.4	48.2	53.3	58.3	60.0	61.9
5	19.4	21.6	23.7	27.0	30.0	33.4	37.5	42.3	47.6	53.4	58.2	60.3	60.8
6	19.1	21.3	24.2	26.8	30.3	34.0	38.0	43.1	48.2	53.4	57.5	61.3	61.5
7	19.3	21.6	23.7	27.1	30.6	33.3	37.6	43.1	48.2	53.2	58.3	60.7	62.2
8	19.3	21.8	24.5	27.1	30.6	33.6	38.4	43.4	47.6	53.6	57.7	60.0	61.9
9	19.0	21.6	24.1	27.0	30.6	34.4	37.7	43.7	49.1	53.6	58.9	60.4	62.5
10	18.9	21.5	24.2	27.4	30.4	33.9	38.7	44.6	49.1	54.1	58.7	60.7	62.3
11	19.2	21.6	24.3	27.3	30.5	34.1	38.6	44.9	48.8	54.0	58.0	60.8	61.9
12	19.2	21.3	24.2	27.5	30.4	35.0	38.6	44.3	49.4	54.3	59.0	61.3	62.3
13	19.1	21.9	24.0	27.4	30.6	34.8	37.6	45.1	49.5	54.6	59.5	60.9	62.6
14	19.1	21.8	24.4	27.6	30.2	34.5	39.0	44.1	50.1	54.7	59.7	61.4	63.2
15	19.5	21.7	24.0	27.6	30.5	34.3	38.3	45.0	49.6	55.3	60.4	61.5	63.5
16	18.9	21.5	24.2	27.6	30.7	34.0	38.8	44.2	49.5	54.6	60.1	60.8	62.3
17	19.1	21.6	24.4	27.2	30.6	34.9	38.7	43.2	49.4	54.5	60.2	61.5	63.4
18	19.0	21.4	24.2	27.2	30.6	34.1	38.1	44.3	49.2	54.3	59.9	62.2	64.8
19	19.0	21.6	24.2	27.2	31.0	34.2	38.1	44.4	49.6	53.6	58.3	62.4	63.3
20	18.9	21.6	23.9	27.1	30.7	34.5	38.9	44.2	49.1	54.1	60.4	61.6	62.6
21	18.9	21.2	24.0	26.7	30.0	33.5	37.7	44.1	49.3	54.0	59.9	61.5	61.5
22	18.7	21.4	24.1	27.1	30.2	34.2	38.2	44.2	48.5	54.4	58.6	60.9	63.9
23	18.9	21.3	23.7	27.3	29.9	33.4	37.5	43.5	49.4	54.0	58.5	60.9	63.8
24	19.0	21.2	24.0	27.1	31.0	34.3	37.3	43.4	48.3	54.2	59.6	61.0	62.8
25	18.8	21.6	24.0	27.1	30.0	34.2	37.9	43.6	48.9	53.6	59.3	60.2	63.9
26	18.7	21.5	24.0	26.8	30.5	33.1	38.6	44.2	48.7	53.9	58.4	61.3	62.5
27	18.8	21.4	24.1	26.7	29.7	33.7	37.8	44.2	48.6	54.2	59.2	60.2	63.0
28	18.7	21.3	24.1	26.8	30.7	34.0	38.0	44.4	49.2	54.2	60.1	62.1	63.1
29	18.8	21.3	24.0	27.4	30.3	34.2	38.7	44.1	49.5	53.9	58.6	60.5	63.8
30	18.9	21.3	23.9	27.2	30.3	33.7	38.3	43.8	48.5	53.4	59.7	60.4	62.3
R 1	19.0	21.4	24.0	27.4	30.7	34.1	39.5	45.0	49.5	53.7	58.1	61.4	62.8
2	19.5	21.7	24.6	27.9	30.8	35.5	38.7	44.3	49.4	54.9	58.7	60.3	62.2
3	19.3	21.6	24.2	27.4	30.9	34.8	38.6	44.9	50.0	54.4	59.6	61.0	62.2
4	19.5	21.8	24.3	27.9	31.3	35.1	39.2	44.9	50.9	54.6	60.1	60.6	62.9
5	19.0	21.7	24.6	27.6	31.0	35.2	39.4	45.3	50.5	54.8	58.7	61.7	63.9

女子

(単位 : kg)

区分	5歳	6	7	8	9	10	11	12	13	14	15	16	17
S 23	15.6	17.6	19.2	21.0	23.0	25.2	27.5	31.6	35.1	39.0	44.1	43.6	48.2
4 3	17.5	18.8	21.1	23.5	25.7	29.4	33.5	38.2	42.7	46.5	48.5	50.0	50.5
6 3	18.7	20.5	22.9	25.7	29.3	33.3	37.3	43.1	47.6	49.2	51.7	53.0	52.8
H 1	18.6	20.8	23.3	26.0	28.9	33.5	38.1	42.9	46.8	49.7	51.6	52.4	52.6
2	18.9	20.8	23.3	26.1	29.9	33.1	38.2	43.2	46.7	49.7	52.0	52.7	52.4
3	19.1	21.0	23.1	26.1	29.1	33.5	38.1	43.8	46.8	49.8	52.3	53.3	52.4
4	18.9	21.0	23.4	26.6	29.5	33.8	39.2	43.4	47.2	50.1	52.3	53.3	52.8
5	18.7	20.8	23.4	26.4	30.2	34.1	38.5	44.1	46.7	49.9	51.5	53.6	54.0
6	18.6	20.9	23.3	26.6	29.4	34.3	39.0	44.5	47.0	50.1	52.0	53.2	53.4
7	19.1	20.7	23.5	26.5	29.9	34.2	38.9	44.2	47.5	50.2	52.5	53.3	52.8
8	18.8	20.8	23.2	26.7	30.6	34.6	39.1	44.4	47.6	50.3	52.1	53.4	53.4
9	18.6	20.9	23.4	26.6	30.2	35.1	39.6	44.6	47.5	50.1	52.2	53.9	53.1
10	18.7	21.2	23.9	26.7	30.2	34.8	40.3	45.4	48.5	50.8	52.2	53.4	53.3
11	18.8	20.9	23.7	26.8	30.1	34.2	40.4	44.8	47.9	50.4	52.0	53.3	52.7
12	18.9	21.1	23.7	27.1	30.5	34.7	39.9	44.8	48.2	50.7	51.7	52.6	53.3
13	18.7	21.4	23.7	26.8	30.7	34.5	38.7	44.2	48.2	50.5	52.5	53.6	53.6
14	19.0	20.7	24.0	26.9	30.7	35.0	40.4	44.8	48.4	50.3	52.1	52.8	53.6
15	18.9	21.2	23.7	27.1	30.4	35.0	39.8	45.0	48.2	50.6	52.0	52.4	53.9
16	18.6	21.0	23.8	26.8	29.8	34.8	39.0	44.8	47.8	50.6	52.2	54.0	53.7
17	18.7	21.3	23.7	26.6	30.5	34.7	40.1	44.5	48.2	50.8	52.2	53.1	54.5
18	18.6	21.0	23.4	26.6	30.3	34.0	40.0	44.1	47.8	50.8	53.1	53.7	53.1
19	18.6	21.0	23.7	26.2	30.1	34.5	39.9	44.6	48.4	50.3	52.8	53.7	53.1
20	18.6	20.9	23.2	26.3	30.4	34.6	39.6	44.9	47.7	50.5	52.0	53.6	53.3
21	18.8	21.0	23.2	26.0	30.3	34.0	39.0	44.9	47.4	50.6	52.8	53.0	54.5
22	18.4	21.0	23.6	26.6	30.4	34.5	39.4	44.5	47.4	50.5	53.6	52.7	52.8
23	18.8	20.9	23.6	26.1	30.0	34.2	39.5	43.8	47.7	49.8	52.6	53.4	53.8
24	18.6	21.1	23.5	25.9	30.0	34.6	39.3	44.3	47.2	49.6	52.2	53.6	53.3
25	18.7	20.8	23.7	26.3	29.7	34.3	38.9	44.3	48.0	50.3	52.7	53.5	52.5
26	18.6	21.0	23.4	26.7	30.4	34.6	39.2	44.7	47.4	50.8	51.2	53.9	52.5
27	18.4	21.2	23.4	26.2	29.7	34.3	39.2	44.5	47.4	50.4	51.7	52.3	54.3
28	18.3	20.7	23.5	26.6	29.8	34.4	39.6	44.4	47.8	50.1	52.7	52.5	53.0
29	18.7	21.2	23.3	26.8	29.9	33.8	39.9	44.4	47.7	51.1	51.7	52.2	53.6
30	18.5	21.1	23.5	26.1	30.6	34.4	38.8	44.1	47.9	50.2	51.7	52.9	53.3
R 1	18.6	21.4	23.5	26.8	30.2	34.5	39.9	44.6	47.8	50.0	51.9	54.1	52.9
2	19.1	21.4	23.7	26.9	30.6	34.5	40.0	45.2	48.1	50.6	51.8	53.9	52.8
3	19.0	21.2	24.1	27.2	30.8	34.7	39.9	44.9	47.8	50.2	52.9	52.7	52.9
4	18.6	21.3	23.8	27.0	30.9	35.2	40.4	44.5	47.7	50.5	51.9	52.2	52.8
5	18.8	21.2	24.0	27.2	31.4	35.5	40.1	45.2	48.4	50.4	52.3	53.0	53.3

注1) 下線の部分は、調査実施以来最高を示す。

注2) 令和2年度から5年度は新型コロナウイルス感染症の影響で、他の年度と測定時期が異なっているため、過去の数値と単純比較することはできない。

(別紙7) 肥満傾向児の出現率の推移 (佐賀県)

(単位：%)

区分	5 歳	6	7	8	9	10	11	12	13	14	15	16	17
H 18	2.68	5.13	6.12	7.05	10.47	11.70	8.88	10.01	9.10	7.87	12.42	10.07	10.48
19	2.34	5.58	6.66	6.41	10.99	9.84	9.19	11.43	10.32	9.24	10.62	11.84	9.51
20	2.27	5.03	6.06	6.85	8.81	11.64	12.23	10.99	8.97	7.94	12.98	9.36	8.38
21	3.33	3.95	4.50	7.16	8.76	7.16	9.90	11.17	9.45	9.43	11.72	9.32	9.81
22	2.18	4.39	5.13	7.97	8.35	10.08	10.45	11.14	8.76	8.50	11.21	9.29	8.32
23	1.85	4.14	5.22	5.79	6.18	8.98	8.50	9.36	8.88	8.06	9.81	8.47	11.08
24	2.21	3.60	6.13	6.81	9.76	9.34	8.51	9.68	7.79	7.10	11.86	9.39	8.23
25	3.04	3.96	5.29	5.48	8.33	10.89	9.90	10.79	9.12	6.88	10.88	6.56	9.86
26	2.15	3.78	5.68	6.60	9.05	8.02	11.78	11.16	9.52	8.49	7.48	11.51	8.13
27	1.45	5.56	5.45	6.88	6.54	8.63	7.83	11.11	7.58	8.22	9.96	6.25	10.10
28	2.58	4.08	4.99	6.06	9.44	9.42	9.91	9.95	9.41	7.74	12.33	7.56	7.23
29	1.79	5.17	5.80	8.59	9.00	9.08	11.16	10.58	10.13	9.01	8.09	7.80	7.68
30	2.51	5.61	7.50	6.12	9.68	11.45	10.43	10.39	8.99	7.90	10.59	6.99	7.50
R 1	3.07	4.99	5.86	9.66	9.46	9.23	12.59	13.16	10.15	8.33	9.73	10.86	7.73
2	3.34	4.28	7.06	9.71	11.99	12.89	10.20	13.36	9.58	10.18	10.03	10.76	11.46
3	3.64	4.30	6.60	9.14	10.53	10.92	10.85	11.92	10.42	8.34	13.40	8.09	7.94
4	2.64	6.20	6.67	9.45	11.85	10.99	12.10	10.51	10.94	9.88	11.60	8.25	9.75
5	1.37	5.87	7.39	8.68	11.29	11.59	10.86	12.34	11.14	10.30	11.01	10.20	11.48

注1) 性別・年齢層・身長別標準体重を求め、肥満度が20%以上の者を肥満傾向児としている。

注2) 令和2年度から5年度は新型コロナウイルス感染症の影響で、他の年度と測定時期が異なっているため、過去の数値と単純比較することはできない。

(別紙8) 痩身傾向児の出現率の推移 (佐賀県)

(単位：%)

区分	5 歳	6	7	8	9	10	11	12	13	14	15	16	17
H 18	0.17	0.47	0.55	0.88	1.73	2.65	2.06	2.39	2.56	1.82	1.19	1.58	0.67
19	0.27	0.96	0.56	0.83	1.17	1.83	2.35	1.99	2.20	1.66	2.00	1.45	1.83
20	0.59	0.69	0.81	0.94	0.68	1.75	1.68	2.09	2.48	2.12	2.28	0.69	0.97
21	0.22	0.21	0.78	1.52	1.43	1.66	3.50	2.88	2.07	1.75	1.87	1.66	1.03
22	0.08	0.65	0.27	0.48	1.12	1.86	2.58	2.21	2.24	1.71	2.26	2.16	0.97
23	-	0.30	0.22	0.22	0.94	1.95	2.11	2.03	1.80	1.45	2.70	1.88	1.27
24	0.80	0.30	0.41	1.26	1.79	1.50	3.08	2.20	1.74	1.94	2.03	1.08	0.83
25	0.47	0.48	0.40	0.80	1.34	1.69	2.50	2.40	1.59	1.49	2.65	1.30	1.29
26	0.10	0.07	0.50	0.99	2.26	2.07	2.97	1.63	2.14	1.79	2.12	1.33	0.71
27	0.33	0.79	0.47	1.00	2.96	1.86	1.64	2.44	2.43	1.19	2.34	1.71	1.55
28	0.39	0.19	0.10	0.10	0.73	1.91	2.81	2.35	1.72	1.48	1.54	2.13	1.07
29	0.37	0.18	0.49	0.69	1.10	1.99	2.53	2.68	2.42	1.72	2.70	1.76	0.89
30	0.25	0.31	0.82	0.49	0.98	2.04	3.15	2.70	2.36	2.04	1.94	1.53	1.12
R 1	0.92	0.44	0.65	1.58	1.42	2.04	2.11	2.14	1.65	1.96	1.82	1.40	0.89
2	0.38	0.35	0.37	0.50	1.37	1.91	3.00	2.68	2.07	2.10	2.57	1.61	1.85
3	0.21	0.28	0.30	0.73	0.71	2.29	2.43	1.78	1.63	2.07	1.96	2.13	1.86
4	0.08	0.23	0.42	1.35	0.75	1.46	2.92	2.78	2.11	1.74	1.68	1.32	1.52
5	0.24	0.21	0.20	0.41	1.50	2.72	3.03	3.03	1.66	1.81	2.45	1.45	2.17

注1) 「-」は、該当者なし

注2) 性別・年齢層・身長別標準体重を求め、肥満度が-20%以下の者を痩身傾向児としている。

注3) 令和2年度から5年度は新型コロナウイルス感染症の影響で、他の年度と測定時期が異なっているため、過去の数値と単純比較することはできない。