

早期水稻定期調査結果(令和6年6月)

25株調査

調査月日：6月14日																
調査地点		葉いもち		紋枯病		セジロウンカ			トビイロウンカ			ヒメビウカ			備考	
		発生株率	発病程度	発生株率	発病程度	発生株率	最高頭数		発生株率	最高頭数		発生株率	最高頭数			
							成虫	幼虫		成虫	幼虫		成虫	幼虫		
A地区	①	0		0		0			0			0				
"	②	0		0		0			0			0				
"	③	0		0		0			0			0				スミリコガイ食害
"	④	0		0		0			0			0				スミリコガイ食害
B地区	①	0		0		4		1 若	0			0				5月調査から圃場変更
"	②	0		0		0			0			0				
"	③	0		0		0			0			0				
"	④	0		0		0			0			0				
C地区	①	0		0		0			0			0				
"	②	0		0		0			0			0				
"	③	0		0		0			0			0				
"	④	0		0		4		1 若	0			0				
発生圃場率		0		0		16.7			0			0				
平均発生率		0		0		0.7			0			0				
発生率の平年値		1.7		0		8.2			0			8.4				
(平年比)		やや少		並		やや少			並			やや少				
発生率の前年値		1.0		0		1.3			0			26.0				