

早植え水稻定期調査結果(令和8年8月上旬)

25株調査

調査月日：8月3日																						
調査地点		品種	葉いもち		紋枯病		セジロウンカ			トビイロウンカ					ヒメトビウンカ			コブノメイガ		フタバコヤカ	備考	
			発生株率	発病程度	発生株率	発病程度	寄生株率	最高頭数 成虫	幼虫 頭	幼虫の齢	寄生株率	最高頭数 成虫	幼虫 頭	幼虫の齢	短翅雌成虫	寄生株率	最高頭数 成虫	幼虫 頭	発生株率	被害程度		寄生株率
D地区	①		%		%		%			%					%			%		%		
	②		0		0		0			0					0			16		0		
	③		0		0		0			0					0			8		0		
	④		4	少	0		0			0					0			40		0		
	④		4	少	0		0			0					0			16		0		
E地区	①		4	少	0		0			0					0			8		0		
	②		12	少	0		8		1	老	0				0			4		0		
	③		0		0		0			0					0			0		0		
	④		12	少	0		4		1	老	0				0			16		0		
F地区	①	にじのきらめき	4	少	0		8		1	老	0				0			4		0		
	②	夢しずく	4	少	0		0			0					0			4		0		
	③	コシヒカリ	0		0		8	1		0					0			8		0		
	④	コシヒカリ	0		0		8	1		0					0			4		4		
発生圃場率			58.3		0		41.7			0					0			91.7		8.3		
平均発生率			3.7		0		3.0			0					0			10.7		0.3		
発生率の平年値			2.1		0.4		3.9			1.6					4.4			2.5		3.1		
（平年比）			並		やや少		並			並					やや少			多		やや少		
発生率の前年値			1.0		0		0			0					0			0		0		